

Pseudo-Goldstini in Field Theory

Riccardo Argurio,^(a) Zohar Komargodski,^(b) and Alberto Mariotti^(c)

^(a) Physique Théorique et Mathématique and International Solvay Institutes
Université Libre de Bruxelles, C.P. 231, 1050 Bruxelles, Belgium

^(b) School of Natural Sciences, Institute for Advanced Study
Einstein Drive, Princeton, NJ 08540, USA

^(c) Theoretische Natuurkunde and International Solvay Institutes
Vrije Universiteit Brussels, Pleinlaan 2, B-1050 Brussels, Belgium

We consider two SUSY-breaking hidden sectors which decouple when their respective couplings to the visible particles are switched off. In such a scenario one expects to find two light fermions: the Goldstino and the pseudo-Goldstino. While the former remains massless in the rigid limit, the latter becomes massive due to radiative effects which we analyze from several different points of view. This analysis is greatly facilitated by a version of the Goldberger-Treiman relation, which allows us to write a universal non-perturbative formula for the mass. We carry out the analysis in detail in the context of gauge mediation, where we find that the pseudo-Goldstino mass is at least around the GeV scale and can be easily at the electroweak range, even in low scale models. This leads to interesting and unconventional possibilities in collider physics and it also has potential applications in cosmology.

1. Introduction

In this note we consider models with multiple supersymmetry-breaking sectors. We assume these SUSY-breaking sectors communicate only through their respective couplings to the Supersymmetric Standard Model (SSM). In other words, the SUSY-breaking sectors decouple when their respective couplings to the SSM are set to zero. Such models could naturally appear in string theory, where there may be several independent sources of supersymmetry breaking. They may also arise naturally in the study of quiver gauge theories. Our main objective is to study the various field-theoretic effects that are relevant in such a setup.

One may wonder whether having such SUSY-breaking sectors which interact only indirectly through the SSM is natural. Indeed, in field theory this can be perfectly natural since renormalizable contact terms may be forbidden by gauge invariance or global symmetries. (By contrast, decoupling in supergravity is a much more delicate question that we will not say anything new about.)

At zeroth order in the interactions with the SSM, there are obviously many massless Goldstini particles. Turning on the small couplings to the SSM, one linear combination, the true Goldstino, remains massless, while the other linear combinations get masses from tree-level and radiative effects.

We will use several methods to analyze these corrections. We first study the problem using the universal chiral Lagrangian for spontaneously broken supersymmetry. The chiral Lagrangian approach shows that the contribution from deep low momenta is quadratically sensitive to the cutoff Λ_{UV} of the chiral Lagrangian

$$m_{G'} \sim \frac{1}{16\pi^2} \frac{m_{gaugino}^3}{f^2} \Lambda_{UV}^2 . \quad (1.1)$$

Hence, the contribution is not dominated by parameterically small momenta and one has to invoke the detailed microscopic physics to determine the mass. The chiral Lagrangian also shows that (1.1) dominates over tree-level contributions that arise due to electroweak symmetry breaking.

As an example of a microscopically well-defined setup we analyze in detail two hidden sectors which only communicate with the SSM via gauge interactions. In this case we find (to all orders in the hidden sector but to leading order in the gauge coupling)

$$m_{G'} = \frac{g^4}{2} \left(\frac{1}{(f^A)^2} + \frac{1}{(f^B)^2} \right) \int \frac{d^4 p}{(2\pi)^4} B_{1/2}^A(p^2) \left(C_0^B - 4C_{1/2}^B + 3C_1^B \right) (p^2) + A \leftrightarrow B , \quad (1.2)$$

where $B_{1/2}^{A,B}, C_0^{A,B}, C_{1/2}^{A,B}, C_1^{A,B}$ are defined through two-point correlation functions of the linear multiplets associated to the hidden sectors A, B . These functions coincide with the functions defined in General Gauge Mediation (GGM) [1]. In order to show that (1.2) is indeed correct, and that the pseudo-Goldstino mass only depends on the functions that appear in GGM, we derive a generalized version of the Goldberger-Treiman relation.

If the two sectors have a common messenger scale and comparable SUSY-breaking scales, one can roughly estimate (1.2) as ~ 1 GeV. On the other hand, we may consider, for instance, different SUSY-breaking scales for the two sectors, then (1.2) can be easily as high as ~ 100 GeV.

We would like to elaborate more on the regime of validity of our analysis. From (1.2) it follows that our field theory effects surely dominate over gravity as long as $m_{3/2} \sim F/M_{PL}$ is smaller than a GeV or so. This means $\sqrt{f} \leq 10^9$, which covers in entirety the parameter space of models based on gauge mediation and variations thereof. On the other hand, since the field theoretic effects can be easily as large as 100 GeV, it may in fact be important to take them into account even in the regime of gravity mediation.

Having such heavy Goldstino-like particles in controllable low scale models potentially leads to unconventional signatures in collider physics and cosmology. Decays of SSM particles sometimes proceed predominantly into the pseudo-Goldstino and may or may not be accompanied by displaced vertices. In addition, the pseudo-Goldstino has three-body decays with observationally interesting time scales.

A recent inspiring paper [2] (for earlier literature on the subject see [3]) considers situations where the gravitational effects are significant. Consistency of SUGRA Lagrangians demands the existence of universal non-renormalizable contact terms mixing the various sectors. Assuming that this is the only source for mixing between the sectors, the authors of [2] computed the supergravity contribution to the mass of the pseudo-Goldstino. They found that the induced mass is $2m_{3/2}$. Possible corrections to this result have been studied in [4] and various interesting applications and variations of this scenario are discussed in [5,6,7]. In this note we consider theories in the rigid limit, where these supergravity corrections are negligible.

The outline of our note is as follows. In section 2 we define more precisely the setup, briefly review some necessary background material about chiral Lagrangians, and discuss in detail contributions to the mass from the deep IR. In section 3 we focus on the scenario where the hidden sectors only communicate with the visible sector via gauge interactions, and present the derivation of (1.2). In section 4 we comment on possible applications to collider physics and cosmology. Two appendices contain technical details that complement the main discussion.

2. Contributions to the Pseudo-Goldstino Mass from Low Energies

2.1. Setup and Review

For simplicity, and without much loss of generality, we will henceforth restrict ourselves to two separate hidden sectors, each communicating with the visible sector in some unspecified way. These interactions with the visible sector could be like in gauge mediation, they can include gauge messengers, Yukawa interactions with SSM fields, some mechanism responsible for $\mu - B_\mu$, and so on.

The two hidden sectors *decouple* if the interactions with the SSM are switched off. Since we neglect supergravity considerations, this form of decoupling can easily be rendered natural due to selection rules imposed by gauge and global symmetries. (On the other hand, decoupling in the full supergravity theory may be more problematic due to moduli; for some general arguments see, for instance, [8,9].) Therefore, our zeroth order problem consists of two utterly decoupled rigid SUSY-breaking theories. Each one of them leads to a massless Goldstino. Let us now review a way of describing the Goldstino theory.

The couplings at low energy of the Goldstino to itself and to other possibly light fields are governed by demanding invariance under non-linearly realized supersymmetry. In general, to write such Lagrangians we only need to insure on-shell invariance under non-linear supersymmetry transformations. However, it is much more convenient to have an off-shell formalism. For this reason we will now briefly review the approach of [10,11]. (There is an extensive list of alternative approaches, e.g. [12] presents several points of view and in [13-18] one can find later discussions. In addition, there are different applications of these ideas, for instance, [19-26] is a sample of some relatively recent activity.)

Consider a general chiral superfield $\overline{D}_{\dot{\alpha}}X = 0$. The superfield X includes a complex scalar, Weyl fermion, and a complex auxiliary field. We can impose an equation

$$X^2 = 0 . \quad (2.1)$$

The (nontrivial) solution is denoted X_{NL} and it is given by

$$X_{NL} = \frac{G^2}{2F} + \sqrt{2}G\theta + \theta^2 F . \quad (2.2)$$

Here the fields are functions of $y = x + i\theta\sigma\bar{\theta}$.¹ Since X_{NL} is a chiral superfield, it can be used to write arbitrary off-shell supersymmetric actions via the usual $\mathcal{N} = 1$ superspace. The simplest possible theory is

$$\mathcal{L} = \int d^4\theta X_{NL}^\dagger X_{NL} + \left(\int d^2\theta f X_{NL} + c.c \right) . \quad (2.3)$$

¹ We are using the conventions of [27] everywhere in this note.

Due to (2.1), this is also the unique theory without (superspace) derivatives. (This has been explicitly shown to be equivalent to the Akulov-Volkov theory. See for example [28-30].) This can be easily extended to include couplings to other light fields, whether they come in complete super-multiplets or not.

Let us assume, for simplicity, that by the time we reach the soft scale E_{soft} (say between 100 GeV and a few TeV), except for the Goldstino, no particles from the hidden sector remain. We can read off the leading low energy couplings between SSM superfields and the Goldstino superfield from various interaction terms in superspace. For example, the gaugino soft-term leads to the following couplings between the observable vector multiplet and the Goldstino

$$\mathcal{L} \supset \frac{m_\lambda}{2f} \int d^2\theta X_{NL} W_\alpha^2 + c.c. . \quad (2.4)$$

This Lagrangian is valid at the energy scale E_{soft} , and is cut off at one of the scales of the hidden sector. In models of weakly coupled supersymmetry breaking the cut off would generically be at the mass scale of the sGoldstino. There is a parameterically sizeable window of energies where (2.4) is valid, and higher derivative corrections can be dropped.

Note that the one-Goldstino couplings in (2.4) are not derivative couplings. However, (2.4) is equivalent to the more familiar description of the Goldstino coupling derivatively to the supercurrent $\sim \partial^\mu G^\alpha S_{\mu\alpha}$. This equivalence can be established by performing a change of variables (which is tantamount to using the equations of motion).

2.2. The Chiral Lagrangian

At zeroth order in the interactions with the SSM, we have a theory of two decoupled Goldstini particles originating from the two hidden sectors labeled A and B

$$\mathcal{L} = \int d^4\theta \left(X_{NL}^{A\dagger} X_{NL}^A + X_{NL}^{B\dagger} X_{NL}^B \right) + \left[\int d^2\theta (f^A X_{NL}^A + f^B X_{NL}^B) + c.c. \right] . \quad (2.5)$$

$f^{A,B}$ and $X_{NL}^{A,B}$ are the corresponding SUSY-breaking scales and constrained superfields.² We now switch on the interactions with the SSM. Both Goldstini couple to the SSM through the couplings (2.4) (and other similar couplings). However, before discussing these couplings, one could consider adding to (2.5) the tree-level term

$$\int d^2\theta m X_{NL}^A X_{NL}^B + c.c. . \quad (2.6)$$

² We can take the $f^{A,B}$ to be real with no loss of generality.

Such a term can be generated by integrating-out the high momentum modes (which one drops from the effective action), and in general no symmetries can be used to set (2.6) to zero. Note that (2.6) generates a mass for the pseudo-Goldstino, but the true Goldstino remains massless, as it should. We cannot compute m using the effective theory, it is an unknown input of the (unspecified) microscopic physics. However, this does not necessarily mean that the effective theory is unavailing; we can still learn something about the typical momentum scales in the problem.

Whatever the corrections induced by (2.4) are, they leave the true Goldstino, G , massless while the pseudo-Goldstino, G' , gets a mass through loops of SSM fields. Denote $f_{eff} = \sqrt{(f^A)^2 + (f^B)^2}$, then the physical Goldstino and pseudo-Goldstino are given by

$$f_{eff}G = f^A G^A + f^B G^B \quad f_{eff}G' = -f^B G^A + f^A G^B . \quad (2.7)$$

At leading order in the supersymmetry breaking parameters, the effective action from integrating out the SSM fields is determined by the one-loop effective Kähler potential

$$K = X_{NL}^{A\dagger} X_{NL}^A + X_{NL}^{B\dagger} X_{NL}^B + K_{1loop}(X_{NL}^A, X_{NL}^{A\dagger}, X_{NL}^B, X_{NL}^{B\dagger}) . \quad (2.8)$$

Once K_{1loop} has been computed, one substitutes the F -term VEVs for $X_{NL}^{A,B}$ in order to extract the non-supersymmetric fermionic masses. Observe that in K_{1loop} each X_{NL} appears at most linearly because of (2.1).

The pseudo-Goldstino mass can be extracted from cubic operators of the form $X_{NL}^A X_{NL}^B X_{NL}^{A\dagger}$ and other operators alike. More precisely, the pseudo-Goldstino mass is given by

$$m_{G'} = \frac{f_{eff}^2}{f^A f^B} (f^B K_{AB\bar{B}} + f^A K_{AB\bar{A}}) . \quad (2.9)$$

Consider two hidden sectors contributing in some way to the soft gaugino mass (assuming a $U(1)$ vector superfield for notational simplicity)

$$\mathcal{L} = \frac{m_\lambda}{2} \int d^2\theta \left(\frac{\alpha^A}{f^A} X_{NL}^A + \frac{\alpha^B}{f^B} X_{NL}^B \right) W_\alpha^2 . \quad (2.10)$$

Note that $\alpha^A + \alpha^B = 1$ by definition of m_λ .

The coupling (2.10) is non-renormalizable, hence we need a generalization of the one-loop Kähler potential to non-renormalizable theories [31]. In this case the effective Kähler potential is quadratically divergent

$$K_{1loop} = -\frac{1}{16\pi^2} \Lambda_{UV}^2 \log \left(1 + m_\lambda \left(\frac{\alpha^A}{f^A} X_{NL}^A + \frac{\alpha^B}{f^B} X_{NL}^B + c.c. \right) \right) . \quad (2.11)$$

where Λ_{UV} is the momentum cutoff in the loop. The resulting pseudo-Goldstino mass is then³

$$m_{G'} = \frac{\alpha^A \alpha^B}{8\pi^2} \left(\frac{f_{eff}}{f^A f^B} \right)^2 m_\lambda^3 \Lambda_{UV}^2, \quad (2.12)$$

and it is *quadratically divergent*. This quadratic divergence can be swallowed in the renormalization of the counterterm (2.6).

A quadratic divergence is bad news because it clearly signifies lack of theoretical control over the exact answer. (Strictly speaking, also finite answers may be prone to corrections from the UV, but cases where the answer is divergent are more obviously UV sensitive.) In other words, the typical momentum in the loop is parameterically larger than the soft scale and therefore the chiral Lagrangian does not give a universal answer. The low energy chiral Lagrangian therefore merely teaches us that one needs to understand the microscopic physics in much more detail; the universal low energy vertices do not suffice.

One can nevertheless try to estimate (2.12) by thinking of Λ_{UV} as the supersymmetric (messenger) scale M . For models like gauge mediation and for comparable SUSY-breaking scales and $\alpha^A \sim \alpha^B$

$$m_{G'} \sim \frac{m_\lambda^3 M^2}{8\pi^2 f^2} \sim 10 \text{ MeV} \quad (\text{effective theory}). \quad (2.13)$$

(We have taken the gauginos to be at the TeV scale and we have used the fact that there are ~ 10 of them in the SSM.) The estimate (2.13) appears to be by and large independent of the messenger scale. Again, since we do not yet have real control over the typical momentum scale of the virtual particles, the estimate (2.13) is only a heuristic first crack at the problem.

One can check that the deep low energy contributions from the B_μ -term, tree level effects, soft non-holomorphic scalar masses, and A -terms are not as significant as (2.13). Some of these facts are established in appendix A.

³ Of course, this result can also be reproduced by an explicit one-loop computation. To perform such computations correctly one must take into account the term bilinear in the Goldstino (2.2). The contact terms stemming from it have a similar role to the seagull term in electrodynamics; they insure the real Goldstino remains massless.

2.3. Integrating-in the sGoldstini

In this subsection we would like to establish that the cutoff in (2.12) is not the sGoldstino mass, even though this is the natural cutoff of the chiral Lagrangian.

At the scale of the sGoldstino particle there is no reason for there not to be many additional resonances. Our purpose here is not to try and write down the most general effective action at this scale, rather, to show that generally the sGoldstini themselves do not render the contribution (2.12) finite.

To include the sGoldstini we simply retain the bottom components of the X fields as propagating degrees of freedom. The simplest way to model this situation is a Polonyi model with an effective Kähler potential that gives a mass to the sGoldstini:

$$\mathcal{L}_{\text{hidden},i} = \int d^4\theta (X_i X_i^\dagger - \frac{1}{\Lambda_i^2} X_i^2 (X_i^\dagger)^2) + \left(\int d^2\theta f_i X_i + c.c. \right), \quad (2.14)$$

with $X_i = x_i + \sqrt{2}\theta\psi_i + \theta^2 F_i$ and $i = A, B$. We might as well parametrize the effective Kähler potential by the pseudomoduli masses and the SUSY-breaking scales through the relations

$$\frac{1}{\Lambda_i^2} = \frac{m_i^2}{4f_i^2}. \quad (2.15)$$

We will assume that the m_i are (well) above the soft scale.

The couplings of the Goldstini superfields to the gauge sector of the SSM are the ones in (2.10), replacing the nonlinear superfields by the linear ones. We can compute the one-loop corrections to the scalar potential for the sGoldstini and subsequently the one-loop mass generated for the pseudo-Goldstino. At leading order in the supersymmetry breaking scale, the one-loop effective Kähler includes both the Coleman-Weinberg effective potential for x_i and the fermionic masses.

The Coleman-Weinberg potential shifts the classical $\langle x_i \rangle = 0$ minimum to

$$\langle x_i \rangle = -\frac{1}{8\pi^2} \frac{\alpha_i m_\lambda^3}{f_i m_i^2} \Lambda_{UV}^2. \quad (2.16)$$

Expanding around this new minimum we can compute all the Goldstino bilinear terms at one-loop and we get

$$\mathcal{L}_{ferm} = \frac{1}{16\pi^2} m_\lambda^3 \Lambda_{UV}^2 \left(\sum_{ij} \frac{\alpha_i \alpha_j G_i G_j}{f_i f_j} - \sum_i \frac{\alpha_i G_i^2}{f_i^2} \right). \quad (2.17)$$

From this we obtain that the induced mass for the pseudo-Goldstino is exactly as in the previous subsection (2.12). Note that taking into account the shift (2.16) of the vacuum expectation value is crucial for finding one massless Goldstino.

The most important conclusion to draw from this discussion is that the pseudo-Goldstino mass is still quadratically divergent even if we include the dynamics of the sGoldstini. The cut off scale Λ_{UV} is thus around the fundamental scale of the theory and the full microscopic theory should be determined.

In the next section we consider a fully specified microscopic setup, and show that the typical momentum is indeed around the fundamental scale. In fact, contributions from this high scale overwhelm the low energy effects we discussed in this section.

3. Hidden Sectors Communicating with the SSM by Gauge Interactions

The next step in our exploration of the various energy scales is to consider a complete microscopic theory. The setup we opt to focus on is depicted in Fig.1. We consider two SUSY-breaking theories, labeled A and B , which communicate with the SSM via gauge interactions. More precisely, when the SSM gauge couplings are set to zero, the sectors A, B decouple from the SSM (and thus also from each other). These decoupled theories have some global symmetry groups in which the SSM gauge group can be embedded and weakly gauged.




Fig.1: Two SUSY-breaking theories communicating with the SSM via gauge interactions.

In essence, this is the setup of GGM [1], only that the hidden sector is assumed to consist of two decoupled field theories. When the gauge couplings are turned on, the two sectors can communicate by exchanging SSM fields. Obviously, at the zeroth order in the gauge couplings, there are two exactly massless Goldstini fermions. Our goal is to find the leading nonzero contribution in an expansion in the gauge couplings.

The mass matrix for the Goldstini system, defined by $-\frac{1}{2}G^i\mathcal{M}_{ij}G^j$ with a symmetric matrix $\mathcal{M}$, is constrained to have one zero eigenvector corresponding to the true Goldstino. Therefore, the matrix has to be of the form

$$\mathcal{M} = \begin{pmatrix} -\frac{f^B}{f^A}\mathcal{M}_{AB} & \mathcal{M}_{AB} \\ \mathcal{M}_{AB} & -\frac{f^A}{f^B}\mathcal{M}_{AB} \end{pmatrix}. \quad (3.1)$$

Once we have calculated $\mathcal{M}_{AB}$, the mass of the pseudo-Goldstino is determined via

$$m_{G'} = \left(\frac{f^B}{f^A} + \frac{f^A}{f^B} \right) \mathcal{M}_{AB} . \quad (3.2)$$

Our goal is therefore to compute the first nontrivial contribution to $\mathcal{M}_{AB}$ in an expansion in the gauge couplings. The processes contributing to $\mathcal{M}_{AB}$ consist of G^A transforming into G^B via some intermediate hidden sector and SSM fields. By virtue of the discussion in appendix A, we can neglect electroweak symmetry breaking effects. In the zeroth order in the gauge coupling g , we cannot have any intermediate SSM fields and thus $\mathcal{M}_{AB} = 0$. In the next order, g^2 , we are allowed to include one intermediate SSM fields, as schematically depicted in Fig.2. Of course this intermediate field must be the gaugino.




Fig.2: At order g^2 the sectors A and B can communicate via one intermediate SSM gaugino.

This process can be forbidden by messenger parity; a symmetry which sends the vector superfield to minus itself and exchanges all the representations accordingly. This forbids a direct zero momentum correlation function between either of $G^{A,B}$ and a gaugino. In fact, gauge mediation models without messenger parity are often unappealing and we will thus assume messenger parity.

We must consider processes of order g^4 . These allow for two intermediate SSM fields and are thus messenger parity invariant. The intermediate fields must be a gaugino and a gauge field or alternatively a gaugino and a D auxiliary field. This is summarized in Fig.3.




Fig.3: At order g^4 the sectors A and B can communicate via two intermediate SSM fields.

We must also add the diagrams with the gauginos flowing in the opposite direction.

In the absence of any particular detailed knowledge of the hidden sector we must account for the blobs formally. On the other hand, if the theory is specified and it is weakly coupled, the blobs can be computed in perturbation theory. For instance, in Minimal Gauge Mediation (MGM) the blobs are, to leading order, triangles with virtual messenger fields. Therefore, the pseudo-Goldstino obtains a mass due to three-loop corrections.

In the processes of Fig.3 the external Goldstini are at zero momentum. One can therefore interpret the blobs as three-point functions of the supercurrent and two insertions of operators of the linear current multiplet. In other words, the pertinent correlation functions are of the form $\langle S_{\nu\alpha}(x)j_\mu(y)\bar{j}_{\dot{\alpha}}(z)\rangle$, $\langle S_{\nu\alpha}(x)J(y)\bar{j}_{\dot{\alpha}}(z)\rangle$, $\langle S_{\nu\alpha}(x)j_\mu(y)j_\beta(z)\rangle$, $\langle S_{\nu\alpha}(x)J(y)j_\beta(z)\rangle$. For our purposes we need the external state to be a zero-momentum Goldstino, therefore, the correlation functions above should be studied only in the limit of large x (much larger than any other scale in the problem).

In this large x limit the three-point functions above simplify dramatically. The reason is that at very low energies the supercurrent flows to the Goldstino particle $S_{\mu\alpha}^{A,B} \sim f^{A,B}\sigma_{\mu\alpha\dot{\alpha}}\bar{G}^{A,B\dot{\alpha}}$ and therefore the large x limit corresponds to inserting a zero momentum Goldstino in the correlation function. This is the same as acting with the supercharge on the vacuum and thus these three-point functions are related to two-point functions of the form $\langle [\bar{Q}_{\dot{\gamma}}, j_\mu(y)\bar{j}_{\dot{\alpha}}(z)] \rangle$, $\langle [\bar{Q}_{\dot{\gamma}}, J(y)\bar{j}_{\dot{\alpha}}(z)] \rangle$, $\langle [\bar{Q}_{\dot{\gamma}}, j_\mu(y)j_\beta(z)] \rangle$, $\langle [\bar{Q}_{\dot{\gamma}}, J(y)j_\beta(z)] \rangle$. These two-point functions, in turn, appear in the calculations of soft masses in gauge mediation. We adopt notation similar to the one in GGM [1]

$$\begin{aligned} \langle J^{A,B}(p)J(-p)^{A,B} \rangle &= C_0^{A,B}(p^2) , \\ \langle j_\alpha^{A,B}(p)\bar{j}_{\dot{\alpha}}^{A,B}(-p) \rangle &= -\sigma_{\alpha\dot{\alpha}}^\mu p_\mu C_{1/2}^{A,B}(p^2) , \\ \langle j_\mu^{A,B}(p)j_\nu^{A,B}(-p) \rangle &= -(p^2\eta_{\mu\nu} - p_\mu p_\nu) C_1^{A,B}(p^2) , \\ \langle j_\alpha^{A,B}(p)j_\beta^{A,B}(-p) \rangle &= \epsilon_{\alpha\beta} B_{1/2}^{A,B}(p^2) . \end{aligned} \tag{3.3}$$

The discussion above shows that the leading order contribution to the pseudo-Goldstino mass should be captured by the functions in (3.3). A quick way to derive the precise relations between these two-point functions and three-point functions is to start by recalling the effective quadratic action for the vector multiplet

$$\frac{1}{g^2}\mathcal{L}_{eff} = \frac{1}{2}C_0^A D^2 - iC_{1/2}^A \lambda\sigma^\mu\partial_\mu\bar{\lambda} - \frac{1}{4}C_1^A F_{\mu\nu}F^{\mu\nu} - \frac{1}{2}\left(B_{1/2}^A\lambda\lambda + c.c.\right) + A \leftrightarrow B . \tag{3.4}$$

This breaks supersymmetry if $B_{1/2} \neq 0$ and if the C s are not all equal. However, it can be supersymmetrized by adding terms linear in the Goldstino as follows

$$\begin{aligned} \frac{1}{g^2}\mathcal{L}_{eff}^{one-G} &= \frac{1}{\sqrt{2}f^A}\left(C_0^A - C_{1/2}^A\right)G^A\sigma^\mu\partial_\mu\bar{\lambda}D + \frac{i}{\sqrt{2}f^A}\left(C_1^A - C_{1/2}^A\right)G^A\sigma_\nu\partial_\mu\bar{\lambda}F^{\mu\nu} \\ &+ \frac{iB_{1/2}^A}{\sqrt{2}f^A}\left(G^A\lambda D - \frac{i}{2}\lambda\sigma^\mu\bar{\sigma}^\nu G^A F_{\mu\nu}\right) + A \leftrightarrow B . \end{aligned} \tag{3.5}$$

To make the theory fully supersymmetric, in addition to (3.5), we need to add terms bilinear in the Goldstini, and terms with derivatives acting on the Goldstini. In order to compute $\mathcal{M}_{AB}$, (3.5) suffices. The procedure we have invoked here is a supersymmetric reincarnation of the Goldberger-Treiman relation. Supersymmetric versions of the Goldberger-Treiman relation have also been useful in the analysis of [32].

From here to derive the mass of the pseudo-Goldstino we only need to carry out the contractions using the vertices in (3.5). After the dust settles, we find that the leading order contribution to the mass of the pseudo Goldstino is

$$m_{G'} = \frac{g^4}{2} \left(\frac{1}{(f^A)^2} + \frac{1}{(f^B)^2} \right) \int \frac{d^4 p}{(2\pi)^4} B_{1/2}^A \left(C_0^B - 4C_{1/2}^B + 3C_1^B \right) + A \leftrightarrow B. \quad (3.6)$$

Note that the combination of the C functions in the integrand is precisely the one appearing in the formula for the soft scalar mass in gauge mediation. The discussion in [33] shows that $C_0 - 4C_{1/2} + 3C_1$ behaves at most like $1/p^4$ at large momentum and it is also possible to prove that $B_{1/2}$ scales at most like $1/p$ at large momentum.⁴ Consequently, the integral is UV convergent.⁵

The computation above has been greatly simplified by the structure of the matrix (3.1), which allowed us to compute $m_{G'}$ only in terms of $\mathcal{M}_{AB}$. As a consistency check, we can also compute the diagonal elements of the mass matrix $\mathcal{M}_{AA}$ and $\mathcal{M}_{BB}$. In order to do this one must take into account also the corrections to (3.5) quadratic in each of the Goldstini.

We can now estimate (3.6) crudely. Assume both hidden sectors have some typical supersymmetric scale M and the SUSY-breaking scales are f^A, f^B . To leading order in the SUSY-breaking scales we would get

$$m_{G'} \sim \frac{g^4}{(16\pi^2)^3} \left(\frac{f^A}{f^B} + \frac{f^B}{f^A} \right) \left(\frac{f^A}{M} + \frac{f^B}{M} \right). \quad (3.7)$$

⁴ The proof of this statement goes as follows. We start from $\langle Q^2 (J(x)J(0)) \rangle \sim \langle j_\alpha(x)j_\beta(0) \rangle$, which was pointed out in [34]. We can now consider the OPE of $J(x)J(0)$. The unit operator is annihilated by Q^2 . Thus, the first interesting contribution in the OPE $J(x)J(0)$ is of the form $\mathcal{O}/x^{4-\Delta_{\mathcal{O}}}$, where unitarity dictates that $\Delta_{\mathcal{O}} > 1$. Thus $\langle j_\alpha(x)j_\beta(0) \rangle$ behaves at small x like $1/x^{3-\epsilon}$ (with some positive ϵ). This implies that in momentum space $B_{1/2}(p^2)$ scales at most like $1/p^{1+\epsilon}$ at large momentum.

⁵ One can also establish IR convergence along the lines of [35].

If the two SUSY-breaking scales are comparable this leads to the estimate

$$m_{G'} \sim \frac{g^4}{(16\pi^2)^3} \frac{f}{M} \sim \frac{g^2}{(16\pi^2)^2} m_{soft} \sim 1 \text{ GeV} . \quad (3.8)$$

(We have included a factor of $\mathcal{O}(10)$ due to the sum over the gauge sector of the SSM.) Note that this is larger than the estimate we obtained in the effective theory (2.13).

However we can also entertain other possibilities. For instance, consider a situation where the fundamental supersymmetric scales in the two sectors are comparable but the SUSY-breaking scales are different. To be concrete we assume that $f^A \gg f^B$ (the soft parameters thus mostly originate in sector A). In this case, the formula (3.7) predicts an enhancement of $m_{G'}$ by f^A/f^B . This ratio, however, cannot be arbitrarily large because at some point the backreaction of the SSM on the hidden sector B becomes too large and our formalism breaks down.⁶ By computing the sGoldstino VEV in sector B , we can estimate that the backreaction is surely tame for $f^A/f^B \ll 10^3$. (For this estimate we have assumed the mass of the sGoldstini is around $f^{A,B}/M$.) Thus, we can easily imagine the pseudo-Goldstino picking a mass at the electroweak range. Note that such a (perhaps surprisingly) large mass for the pseudo-Goldstino is achieved effortlessly and ubiquitously in low scale models, where corrections from supergravity are completely negligible.

One can also evaluate (3.6) explicitly in a variety of simple realizations of gauge mediation. In appendix B we consider one such example, where the two hidden sectors are copies of MGM.

4. Phenomenology of Goldstini

In the scenario presented in this note, the pseudo-Goldstino is generically the Next-to-Lightest Supersymmetric Particle (NLSP), with the LSP being of course the very light gravitino. The pseudo-Goldstino is not stable and its decay can be analyzed via the chiral Lagrangian. For instance, the terms responsible for the gaugino mass (2.10) give rise to vertices of the form $\sim G\sigma_\mu\bar{\sigma}_\nu\lambda F^{\mu\nu}$ which induce three-body decays of the pseudo-Goldstino into two photons and the true Goldstino. There are also some very important vertices with two Goldstini. In fact, the naive estimate based on dimensional analysis fails due to an exact cancelation between the different vertices. An analogous story takes place in the

⁶ This is similar to the breakdown of GGM if at zero gauge coupling there is no stable vacuum in the hidden sector.

couplings to the SM fermions. One is left with the following estimate of the decay width into two standard model fermions and the true Goldstino [6]

$$\Gamma_{G' \rightarrow G f \bar{f}} \sim \frac{m_{G'}^9}{10^5 f_{eff}^4} \left(\frac{(m_{\tilde{f}}^A)^2 \tan \theta - (m_{\tilde{f}}^B)^2 \cot \theta}{m_{\tilde{f}}^2} \right)^2. \quad (4.1)$$

We denote $\tan \theta = f^B/f^A$ and $(m_{\tilde{f}}^{A,B})^2$ are the contributions to the mass of the slepton from the two hidden sectors, such that $(m_{\tilde{f}}^A)^2 + (m_{\tilde{f}}^B)^2 = m_{\tilde{f}}^2$. There is a similar width to decay into two photons and the true Goldstino.

Consider theories with two general SUSY-breaking scales $f^A \geq f^B$. Assuming again, for simplicity, that the messenger scales in the two sectors are comparable and taking $m_{\tilde{f}}^{A,B} \sim f^{A,B}/M$ we find

$$\tau \sim 10^{21} \text{ sec} \left(\frac{f_{eff}}{10^{10} \text{ GeV}^2} \right)^4 \left(\frac{f^B}{f^A} \right)^7. \quad (4.2)$$

To derive the estimate above we have taken the mass of the pseudo Goldstino to be $m_{G'} \sim f^A/f^B \text{ GeV}$. This gives rise to many different possibilities. For instance, when the pseudo-Goldstino is around the weak or TeV scale (i.e. $f^A/f^B \sim 10^{2-3}$) models of low scale mediation $\sqrt{f} \sim 10^{4-5} \text{ GeV}$ give a lifetime of the order of a few seconds. Still keeping the pseudo-Goldstino at the weak-TeV scale, we can also choose $\sqrt{f_{eff}} \sim 10^8 \text{ GeV}$ which leads to lifetimes of the order of 10^{23-24} secs. Both of these time scales have potentially interesting observable consequences [36]. One can of course imagine many other scenarios stemming from (4.2), including scenarios with lighter pseudo-Goldstino.

One can also easily imagine many unconventional collider manifestations of the setup here.⁷ One obvious consequence of having two different hidden sectors is that the relation between the decay time of the Lightest Observable-sector Supersymmetric Particle (LOSP) and the scale of SUSY breaking is no longer universally determined when the decay is at least for a significant fraction to the pseudo-Goldstino. This can have several different consequences.

For instance, consider two hidden sectors with comparable messenger scales but with a possible hierarchy in the SUSY breaking scales. From the couplings (2.10) we see that the gaugino is equally likely to decay to either of the Goldstini (since the dependence on

⁷ We thank A. Katz for discussions on this topic.

f cancels and only the supersymmetric scale remains). Therefore, if the LOSP is bino- or wino-like, and it is heavier than the pseudo-Goldstino, many of the processes of the SSM will terminate in a heavy, long lived, pseudo-Goldstino (the decay can be prompt or there can be displaced vertices). This also comes accompanied by an isolated photon from the last step of the decay. Having such an invisible heavy particle as missing energy is clearly different from conventional scenarios of gauge mediation where the missing energy is carried away by practically massless objects. It is also distinguishable from gravity mediation, where the LOSP is stable on collider time scales and therefore, if it is a gaugino, no isolated photons are expected.

We will not attempt to classify all the scenarios and signatures here. The very brief remarks above are just to demonstrate that unusual collider and cosmological signatures are definitely possible. Clearly, it will be interesting to investigate the various possibilities further. It is also important to study more general hidden sector paradigms, beyond gauge mediation.

Acknowledgments: We would like to thank M. Bertolini, M. Buican, T. Dumitrescu, G. Ferretti, A. Katz, K. Mawatari, N. Seiberg and D. Shih for helpful conversations. R.A. is a Research Associate of the Fonds de la Recherche Scientifique–F.N.R.S. (Belgium). The research of R.A. is supported in part by IISN-Belgium (conventions 4.4511.06, 4.4505.86 and 4.4514.08) and by the “Communauté Française de Belgique” through the ARC program. A.M. is a Postdoctoral Researcher of FWO-Vlaanderen. A.M. is also supported in part by FWO-Vlaanderen through project G.0114.10N. R.A. and A.M. are supported in part by the Belgian Federal Science Policy Office through the Interuniversity Attraction Pole IAP VI/11. Z.K. gratefully acknowledges support by DOE grant DE-FG02-90ER40542. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation.

Appendix A. Other Contributions from Low Energies

A.1. The B_μ -term

The B_μ -term leads to the following coupling between the observable Higgs multiplets and the Goldstino

$$\mathcal{L} \supset \frac{B_\mu}{f} \int d^2\theta X_{NL} H_u H_d + c.c. \quad . \quad (\text{A.1})$$

In the case at hand, we have two different SUSY-breaking sectors, each contributing independently to the B_μ -term

$$\int d^2\theta B_\mu \left(\frac{\lambda^A}{f^A} X_{NL}^A H_u H_d + \frac{\lambda^B}{f^B} X_{NL}^B H_u H_d \right) + \int d^2\theta \mu H_u H_d + c.c. . \quad (\text{A.2})$$

Of course $\lambda^A + \lambda^B = 1$ since B_μ is the actual physical soft term. We also included the μ -term in (A.2).

The effective Kähler potential from integrating out the SSM Higgs fields (A.2) (which for our purpose appear only quadratically) is [37,38] (we keep only the significant terms)

$$K_{1loop} = -\frac{1}{32\pi^2} \text{Tr} \left[\mathcal{M} \mathcal{M}^\dagger \log \frac{\mathcal{M} \mathcal{M}^\dagger}{\Lambda^2} \right] , \quad (\text{A.3})$$

where

$$\mathcal{M} = B_\mu \begin{pmatrix} 0 & \frac{\lambda^A}{f^A} X_{NL}^A + \frac{\lambda^B}{f^B} X_{NL}^B + \frac{\mu}{B_\mu} \\ \frac{\lambda^A}{f^A} X_{NL}^A + \frac{\lambda^B}{f^B} X_{NL}^B + \frac{\mu}{B_\mu} & 0 \end{pmatrix} . \quad (\text{A.4})$$

The resulting contribution to the mass of the pseudo-Goldstino is

$$m_{G'} = \frac{\lambda^A \lambda^B}{16\pi^2} \left(\frac{f_{eff}}{f^A f^B} \right)^2 \frac{B_\mu^3}{\mu} . \quad (\text{A.5})$$

The result of this calculation in the chiral Lagrangian is finite.

If the B_μ -term is around the soft scale, and SUSY breaking occurs at low scales ($f \sim 10^8 \text{ GeV}^2$), (A.5) is of the order of 10 eV but it can increase to as much as an MeV if B_μ is larger by an order of magnitude. This is numerically smaller than the effect from low momentum vector multiplet loops (2.13). The typical momentum of the virtual particles is low and the result of the low energy calculation is convergent. This however, does not necessarily mean the result (A.5) is reliable; there could still be contributions from the UV.

A.2. Tree-Level Contributions

Effects of electroweak symmetry breaking induce tree-level contributions to the pseudo-Goldstino mass. For instance, due to electroweak symmetry breaking, the hypercharge D -term is generally nonzero. Let us denote its value simply by D . Then, the gaugino soft mass terms (2.10) lead to the following mass matrix in the space of the three fermions G^A, G^B, λ

$$\mathcal{M} = m_\lambda \begin{pmatrix} \frac{\alpha^A D^2}{2(f^A)^2} & 0 & \frac{\alpha^A D}{\sqrt{2}f^A} \\ 0 & \frac{\alpha^B D^2}{2(f^B)^2} & \frac{\alpha^B D}{\sqrt{2}f^B} \\ \frac{\alpha^A D}{\sqrt{2}f^A} & \frac{\alpha^B D}{\sqrt{2}f^B} & 1 \end{pmatrix} . \quad (\text{A.6})$$

If the hypercharge D -term is set to zero, the only tree-level contribution that remains is the gaugino mass term and the pseudo-Goldstino is massless at tree-level. In this case only radiative effects exist. The matrix (A.6) has, schematically (assuming the two SUSY-breaking scales are comparable and neglecting corrections in D/f), the eigenvalues

$$\left(0, \frac{m_\lambda D^2}{f^2}, m_\lambda\right), \quad (\text{A.7})$$

corresponding to the real Goldstino, the pseudo-Goldstino, and the gaugino, respectively. Therefore, the tree-level contribution to the mass of the pseudo-Goldstino scales like

$$m_{G'}^{(tree)} \sim m_\lambda \frac{D^2}{f^2}, \quad (\text{A.8})$$

which is parameterically smaller than the low-energy quantum effect (2.13). For instance, (A.8) is maximized for low scale models ($\sqrt{f} \sim 10$ TeV), where it can be roughly estimated as 10 keV.

An additional tree-level effect could arise from the Higgs superfields picking up an F -term. For instance, the source for the B_μ term (A.2) also induces, upon electroweak symmetry breaking, a tree-level mass for G' . With similar techniques, we can estimate it to be parameterically smaller than (2.13).

Tree-level contributions arising from electroweak breaking are therefore negligible.

Appendix B. Two Copies of MGM

Here we consider an explicit example of the class of models studied in section 3. Suppose that each of the two hidden sectors is a copy of Minimal Gauge Mediation (MGM). For simplicity we discuss the case in which the SSM is represented by a $U(1)$ gauge group. The messengers $\Phi_i, \tilde{\Phi}_i$ are massive and interact with the spurions X^i ($i = A, B$) as

$$\mathcal{L}_i = \int d^2\theta \left(h_i X^i \Phi_i \tilde{\Phi}_i + M_i \Phi_i \tilde{\Phi}_i \right), \quad (\text{B.1})$$

where the spurions X^i acquire F -term VEVs f^i and contain the Goldstini in the θ components. We assume without loss of generality that the bottom components x^i are stabilized at the origin.

The leading contributions to $\mathcal{M}_{AB}$ are easily listed. They consist of triangle diagrams fused by SSM fields. In each of the diagrams, only one of the triangles needs an F -term insertion. We summarize the various three-loop diagrams in Fig.4.




Fig.4: The three-loop diagrams contributing to $\mathcal{M}_{AB}$.

The computation can be done in two steps. First we compute the momentum dependence of the triangle vertices when the Goldstino particle is at zero momentum and the gauge-multiplet field have some general momentum p . To compute the three-loop diagrams we then have to convolve these momentum dependent vertices with the gauge-multiplet propagators and perform the integration. This mimics the procedure we have carried out when discussing the general framework in section 3.

We only display the computations to first order in the supersymmetry breaking scales f^A and f^B . Doing the explicit computation we find that the vertices in momentum space with zero momentum Goldstini are

$$\begin{aligned} \frac{1}{g^2} \mathcal{L}_{eff}^{one-G} &= \frac{iV_0^A(p^2/M_A^2)}{\sqrt{2}f^A} \left(G^A \lambda D - \frac{i}{2} \lambda \sigma^\mu \bar{\sigma}^\nu G^A F_{\mu\nu} \right) \\ &+ \frac{iV_1^A(p^2/M_A^2)}{\sqrt{2}f^A} G^A \sigma^\mu p_\mu \bar{\lambda} D - \frac{V_2^A(p^2/M_A^2)}{\sqrt{2}f^A} G^A \sigma_\nu p_\mu \bar{\lambda} F^{\mu\nu} + A \leftrightarrow B . \end{aligned} \quad (\text{B.2})$$

The functions $V_0^A(p^2/M_A^2)$, $V_1^A(p^2/M_A^2)$, $V_2^A(p^2/M_A^2)$ are given by

$$\begin{aligned} V_0^A(p^2/M_A^2) &= \frac{1}{16\pi^2} \frac{f^A}{M_A} b(p^2/M_A^2) = \frac{1}{16\pi^2} \frac{f_A}{M_A} \int_0^1 dy \frac{1}{1+y(1-y)x_A} , \\ V_1^A(p^2/M_A^2) &= V_2^A(p^2/M_A^2) = \frac{1}{16\pi^2} \frac{f_A^2}{M_A^4} v(p^2/M_A^2) = \frac{1}{16\pi^2} \frac{(f^A)^2}{M_A^4} \int_0^1 dy \frac{y^2(1-y)}{(1+y(1-y)x_A)^2} , \end{aligned} \quad (\text{B.3})$$

where $x_A = p^2/M_A^2$, and the y integrals can be done analytically. Comparing these with the expressions for the B and C functions of MGM (see [1]) one can verify that the momentum dependent vertices are exactly as in (3.5). (Note in particular that at this order in SUSY-breaking for MGM we have $C_1 = C_0$.) The final answer for the pseudo-Goldstino mass is then

$$m_{G'} = \frac{g^4}{(16\pi^2)^3} \left(\frac{1}{(f^A)^2} + \frac{1}{(f^B)^2} \right) \int_0^\infty dx x \frac{f^A (f^B)^2}{M_A} v(x) b\left(\frac{xM_B^2}{M_A^2}\right) + A \leftrightarrow B . \quad (\text{B.4})$$

We can calculate this integral in different limits. For $M_A = M_B \equiv M$ we get

$$m_{G'} \simeq 4.21 \times \frac{g^4}{(16\pi^2)^3} \frac{((f^A)^2 + (f^B)^2)(f^A + f^B)}{f^A f^B M} , \quad (\text{B.5})$$

in agreement with (3.7). We can also consider cases where the messenger masses are not the same. This may open up new interesting possibilities, but we leave it for the future.

References

- [1] P. Meade, N. Seiberg and D. Shih, “General Gauge Mediation,” arXiv:0801.3278 [hep-ph].
- [2] C. Cheung, Y. Nomura and J. Thaler, “Goldstini,” JHEP **1003**, 073 (2010) [arXiv:1002.1967 [hep-ph]];
- [3] K. Benakli, C. Moura, “Brane-Worlds Pseudo-Goldstinos,” Nucl. Phys. **B791**, 125-163 (2008). [arXiv:0706.3127 [hep-th]].
- [4] N. Craig, J. March-Russell and M. McCullough, “The Goldstini Variations,” JHEP **1010**, 095 (2010) [arXiv:1007.1239 [hep-ph]].
- [5] C. Cheung, J. Mardon, Y. Nomura and J. Thaler, “A Definitive Signal of Multiple Supersymmetry Breaking,” JHEP **1007**, 035 (2010) [arXiv:1004.4637 [hep-ph]].
- [6] H. C. Cheng, W. C. Huang, I. Low and A. Menon, “Goldstini as the decaying dark matter,” arXiv:1012.5300 [hep-ph].
- [7] K. -I. Izawa, Y. Nakai, T. Shimomura, “Higgs Portal to Visible Supersymmetry Breaking,” [arXiv:1101.4633 [hep-ph]].
- [8] S. Kachru, J. McGreevy and P. Svrcek, “Bounds on masses of bulk fields in string compactifications,” JHEP **0604**, 023 (2006) [arXiv:hep-th/0601111].
- [9] Z. Komargodski and N. Seiberg, “Comments on Supercurrent Multiplets, Supersymmetric Field Theories and Supergravity,” JHEP **1007**, 017 (2010) [arXiv:1002.2228 [hep-th]].
- [10] R. Casalbuoni, S. De Curtis, D. Dominici, F. Feruglio and R. Gatto, “Nonlinear Realization of Supersymmetry Algebra from Supersymmetric Constraint,” Phys. Lett. B **220**, 569 (1989).
- [11] Z. Komargodski and N. Seiberg, “From Linear SUSY to Constrained Superfields,” arXiv:0907.2441 [hep-th].
- [12] S. Samuel and J. Wess, “A Superfield Formulation Of The Nonlinear Realization Of Supersymmetry And Its Coupling To Supergravity,” Nucl. Phys. B **221**, 153 (1983).
- [13] T. E. Clark and S. T. Love, “Goldstino couplings to matter,” Phys. Rev. D **54**, 5723 (1996) [arXiv:hep-ph/9608243].
- [14] A. Brignole, F. Feruglio and F. Zwirner, “Aspects of spontaneously broken $N = 1$ global supersymmetry in the presence of gauge interactions,” Nucl. Phys. B **501**, 332 (1997) [arXiv:hep-ph/9703286].
- [15] M. A. Luty and E. Ponton, “Effective Lagrangians and light gravitino phenomenology,” Phys. Rev. D **57**, 4167 (1998) [arXiv:hep-ph/9706268].
- [16] A. Brignole, F. Feruglio and F. Zwirner, “On the effective interactions of a light gravitino with matter fermions,” JHEP **9711**, 001 (1997) [arXiv:hep-th/9709111].
- [17] M. Klein, “Couplings in pseudo-supersymmetry,” Phys. Rev. D **66**, 055009 (2002) [arXiv:hep-th/0205300].

- [18] A. Brignole, J. A. Casas, J. R. Espinosa and I. Navarro, “Low-scale supersymmetry breaking: Effective description, electroweak breaking and phenomenology,” Nucl. Phys. B **666**, 105 (2003) [arXiv:hep-ph/0301121].
- [19] I. Antoniadis and M. Tuckmantel, “Non-linear supersymmetry and intersecting D-branes,” Nucl. Phys. B **697**, 3 (2004) [arXiv:hep-th/0406010].
- [20] I. Antoniadis, M. Tuckmantel and F. Zwirner, “Phenomenology of a leptonic goldstino and invisible Higgs boson decays,” Nucl. Phys. B **707**, 215 (2005) [arXiv:hep-ph/0410165].
- [21] M. Dine, R. Kitano, A. Morisse and Y. Shirman, “Moduli decays and gravitinos,” Phys. Rev. D **73**, 123518 (2006) [arXiv:hep-ph/0604140].
- [22] M. L. Graesser, R. Kitano and M. Kurachi, “Higgsinoless Supersymmetry and Hidden Gravity,” JHEP **0910**, 077 (2009) [arXiv:0907.2988 [hep-ph]].
- [23] M. Dine, G. Festuccia and Z. Komargodski, “A Bound on the Superpotential,” JHEP **1003**, 011 (2010) [arXiv:0910.2527 [hep-th]].
- [24] L. Alvarez-Gaume, C. Gomez and R. Jimenez, “Minimal Inflation,” Phys. Lett. B **690**, 68 (2010) [arXiv:1001.0010 [hep-th]].
- [25] I. Antoniadis and M. Buican, “Goldstinos, Supercurrents and Metastable SUSY Breaking in N=2 Supersymmetric Gauge Theories,” arXiv:1005.3012 [hep-th].
- [26] I. Antoniadis, E. Dudas, D. M. Ghilencea and P. Tziveloglou, “Non-linear MSSM,” Nucl. Phys. B **841**, 157 (2010) [arXiv:1006.1662 [hep-ph]].
- [27] J. Wess and J. Bagger, “Supersymmetry and supergravity,” *Princeton, USA: Univ. Pr. (1992) 259 p*
- [28] H. Liu, H. Luo, M. Luo and L. Wang, “Leading Order Actions of Goldstino Fields,” arXiv:1005.0231 [hep-th].
- [29] A. A. Zheltukhin, “On equivalence of the Komargodski-Seiberg action to the Volkov-Akulov action,” arXiv:1009.2166 [hep-th].
- [30] S. M. Kuzenko and S. J. Tyler, “Relating the Komargodski-Seiberg and Akulov-Volkov actions: Exact nonlinear field redefinition,” arXiv:1009.3298 [hep-th].
- [31] A. Brignole, “One-loop Kaehler potential in non-renormalizable theories,” Nucl. Phys. B **579**, 101 (2000) [arXiv:hep-th/0001121].
- [32] J. A. Bagger and A. F. Falk, “Decoupling and Destabilizing in Spontaneously Broken Supersymmetry,” Phys. Rev. D **76**, 105026 (2007) [arXiv:0708.3364 [hep-ph]].
- [33] T. T. Dumitrescu, Z. Komargodski, N. Seiberg and D. Shih, “General Messenger Gauge Mediation,” JHEP **1005**, 096 (2010) [arXiv:1003.2661 [hep-ph]].
- [34] M. Buican, P. Meade, N. Seiberg and D. Shih, “Exploring General Gauge Mediation,” JHEP **0903**, 016 (2009) [arXiv:0812.3668 [hep-ph]].
- [35] M. Buican and Z. Komargodski, “Soft Terms from Broken Symmetries,” JHEP **1002**, 005 (2010) [arXiv:0909.4824 [hep-ph]].

- [36] A. Arvanitaki, S. Dimopoulos, S. Dubovsky et al., “Astrophysical Probes of Unification,” *Phys. Rev.* **D79**, 105022 (2009). [arXiv:0812.2075 [hep-ph]].
- [37] I. L. Buchbinder, S. Kuzenko, Z. Yarevskaya, “Supersymmetric effective potential: Superfield approach,” *Nucl. Phys.* **B411**, 665-692 (1994).
- [38] M. T. Grisaru, M. Rocek and R. von Unge, “Effective Kähler Potentials,” *Phys. Lett. B* **383**, 415 (1996) [arXiv:hep-th/9605149].